



## Pinnacle Orthopaedics Hip Injection Discharge Instructions

### GENERAL ANESTHESIA OR SEDATION

- Do not drive or operate machinery for at least 24 hours
- Do not consume alcohol, tranquilizers, sleeping medications, or any non prescribed medication for at least 48 hours
- Do not make important decisions or sign any important papers for at least the next 48 hours
- You should have someone with you at home tonight. Children may appear flushed for several hours after surgery

### ACTIVITY

- You are advised to go directly home from the hospital. Restrict your activities and rest for the day. Resume light to normal activities tomorrow.
- Do not engage in strenuous activity that may place stress on your incision
- You have **NO** motion or weight bearing restrictions and can start working on range of motion without concern.

### FLUIDS AND DIET

- Begin with clear liquids, bullion, dry toast, soda crackers. If not nauseated, you may go to a regular diet when you desire.
- Greasy and spicy foods are not advised for today.

### MEDICATIONS

- Take over the counter medications such as Tylenol, Motrin, or aleve (unless contra-indicated) to help reduce postoperative pain and swelling.
- You may resume your daily prescription medication schedule.

### OPERATIVE SITE/ PROCEDURE

- Keep dressing clean and dry for 1-2 days
- You May wash over the injection site with soap and water in the shower in 1-2 days.

### EXTREMITIES: ARMS, HANDS, LEGS, FEET

- Keep operative extremity elevated as much as possible to lessen swelling and discomfort.
- Full weight bearing on operative extremity.

### NORMAL SIGNS/ SYMPTOMS POST PROCEDURE

- "Feelings" of tiredness / mild disorientation
- "Feelings" of nausea or lack of appetite may also experience some minor episodes of vomiting
- "Feelings" of soreness/discomfort/pain, which should be relieved by medication
- "Feelings" of numbness or tingling may last for an acceptable period, depending on your surgical procedure
- "Normal" for an operative site to appear slightly swollen and pink or have small areas of drainage. .

### FOLLOW-UP CARE: SPECIFICALLY WATCH FOR:

- Fever over 101 degrees by mouth.
- Increased redness, warmth, hardness and around operative area
- Inability to urinate
- Blood soaked dressing
- Swelling around operative area
- Increasing and progressive draining from surgical area or exam site
- Pain not relieved by medication
- Numbness, tingling, cold fingers or toes
- Persistent nausea and/or vomiting; unable to retain fluids

**Call your Pinnacle Orthopaedics if you have any problems that concern you.**

**IF YOU NEED IMMEDIATE ATTENTION, GO TO THE NEAREST EMERGENCY ROOM OR CALL 911**

An appointment for your post-op checkup has been made for: \_\_\_\_\_

I want to see you back in 2 weeks

I have received a copy of these discharge instructions:

Patient/Responsible adult signature \_\_\_\_\_